

PHARMACOPOEIAL DISCUSSION GROUP

CORRECTION

CODE: E-49

NAME: PROPYL PARAHYDROXYBENZOATE

(Correction 2 to revision 1 of the sign-off document signed 10 June 2009)

Item to be corrected:

- Addition of CAS numbers: [94-13-3]
- Appearance of solution/color: addition of comparison with alcohol

Attribute	EP	JP	USP
Definition	+	+	+
Identification A (melting point)*	+	+	+
Identification B (IR)	+	+	+
Appearance of solution/color	+	+	+
Acidity	+	+	+
Related substances**	+	+	+
Sulphated ash	+	+	+
Assay	+	+	+

* Melting point: listed in JP as a test and not as part of identification

** Related substances: JP uses the term "relative response factor" instead of "correction factor"

Legend

+ will adopt and implement

- will not stipulate

Non-harmonised attributes

Characters, Storage


Local requirements

Ph. Eur.	JP	USP
Second identification (melting point, TLC)	Related substances: test for required detectability, system repeatability Heavy metals (20 ppm) Assay: column temperature	none

Reagents and reference materials

Each pharmacopoeia will adapt the text to take account of local reference materials and reagent specifications.


European Pharmacopoeia

Signature 

Name
VIGOR GAMB

Date
22-Dec-2020


Japanese Pharmacopoeia

Signature

for Y. Yoshida

Name
Humberto Olveda

Date
16 Dec/2020


United States Pharmacopoeia

Signature


Name
KEVIN MASEC

Date
19-NOV-2020

E49 - PROPYL PARAHYDROXYBENZOATE

C₁₀H₁₂O₃

Mr 180.2

[94-13-3]

DEFINITION

Propyl 4-hydroxybenzoate.

Content: 98.0 per cent to 102.0 per cent.

IDENTIFICATION

A. *Melting point:* 96 °C to 99 °C.

B. *Infrared absorption spectrophotometry.*

Record the infrared absorption spectrum of propyl parahydroxybenzoate and compare with the Reference Spectrum or the spectrum obtained with the Reference Standard: the transmission minima correspond in position and relative size.

TESTS

Solution S. Dissolve 1.0 g in *alcohol* and dilute to 10 ml with the same solvent.

Appearance of solution. Solution S is clear and not more intensely coloured than *alcohol* or the reference solution.

Primary solutions:

- *Ferric chloride primary solution:* a 45.0 g/l solution of ferric chloride (FeCl₃, 6H₂O).
- *Cobalt chloride primary solution:* a 59.5 g/l solution of cobalt chloride (CoCl₂, 6H₂O).
- *Copper sulphate primary solution:* a 62.4 g/l solution of copper sulphate (CuSO₄, 5H₂O).

Reference solution:

To 5.0 ml of cobalt chloride primary solution, 12.0 ml of ferric chloride primary solution and 2.0 ml of copper sulphate primary solution, add hydrochloric acid (10 g/l HCl) to make 1000.0 ml.

Acidity. To 2 ml of solution S add 3 ml of *alcohol*, 5 ml of *carbon dioxide-free water* and 0.1 ml of *bromocresol green solution*. Not more than 0.1 ml of 0.1 M *sodium hydroxide* is required to change the colour of the indicator to blue.

Related substances. Liquid chromatography.

Test solution. Dissolve 50.0 mg of the sample to be examined in 2.5 ml of *methanol* and dilute to 50.0 ml with the mobile phase. Dilute 10.0 ml of this solution to 100.0 ml with the mobile phase.

Reference solution (a). Dissolve 5 mg each of *ethyl parahydroxybenzoate R*, *4-hydroxybenzoic acid R* and the substance to be examined in the mobile phase and dilute to 100.0 ml with the same solvent. Dilute 1 ml of this solution to 10.0 ml with the mobile phase.

Reference solution (b). Dissolve 50.0 mg of *propyl parahydroxybenzoate CRS* in 2.5 ml of *methanol* and dilute to 50.0 ml with the mobile phase. Dilute 10.0 ml of this solution to 100.0 ml with the mobile phase.

Reference solution (c). Dilute 1.0 ml of the test solution to 20.0 ml with the mobile phase. Dilute 1.0 ml of this solution to 10.0 ml with the mobile phase.

Column:

— size: $l = 0.15$ m, $\varnothing = 4.6$ mm;

— stationary phase: *octadecylsilyl silica gel for chromatography* (5 μm).

Mobile phase: 6.8 g/l solution of *potassium dihydrogen phosphate*, *methanol* (35:65 V/V).

Flow rate: 1.3 ml/min.

Detection: 272 nm.

Injection: 10 μl of the test solution and reference solutions (a) and (c).

Run time: 2.5 times the retention time of *propyl parahydroxybenzoate*.

Relative retention with reference to *propyl parahydroxybenzoate* (retention time = about 4.5 min):
4-hydroxybenzoic acid = about 0.3; *ethyl parahydroxybenzoate* = about 0.7.

System suitability:

- *resolution:* minimum of 3.0 between the peaks due to *ethyl parahydroxybenzoate* and to *propyl parahydroxybenzoate* in the chromatogram obtained with reference solution (a).

Limits:

- *correction factor:* for the calculation of content, multiply the peak area of *4-hydroxybenzoic acid* by 1.4;

- *4-hydroxybenzoic acid:* not more than the area of the principal peak in the chromatogram obtained with reference solution (c) (0.5 per cent);

- *unspecified impurities:* for each impurity, not more than the area of the principal peak in the chromatogram obtained with reference solution (c) (0.5 per cent);

- *total:* not more than twice the area of the principal peak in the chromatogram obtained with reference solution (c) (1.0 per cent);

- *disregard limit:* 0.2 times the area of the principal peak in the chromatogram obtained with reference solution (c) (0.1 per cent).

Sulphated ash: maximum 0.1 per cent, determined on 1.0 g.

ASSAY

Liquid chromatography as described in the test for related substances with the following modification.

Injection: test solution and reference solution (b).

System suitability:

- *repeatability:* maximum relative standard deviation of 0.85 per cent after 6 injections of the reference solution (b).

Calculate the percentage content of $C_{10}H_{12}O_3$ in the sample to be examined from the peak areas in the chromatograms obtained with test solution and reference solution (b) and the declared content of *propyl parahydroxybenzoate CRS*.

REAGENTS

Bromocresol green solution.

Dissolve 50 mg of *bromocresol green* in 0.72 ml of 0.1 M *sodium hydroxide* and 20 ml of *alcohol* and dilute to 100 ml with *water*.

Test for sensitivity. To 0.2 ml of the bromocresol green solution add 100 ml of *carbon dioxide-free water*. The solution is blue. Not more than 0.2 ml of 0.02 M *hydrochloric acid* is required to change the colour to yellow.

Colour change: pH 3.6 (yellow) to pH 5.2 (blue).